

Kin Conflict in Seed Development: An Interdependent but Fractious Collective

David Haig

Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts 02138; email: dhaig@oeb.harvard.edu

Annu. Rev. Cell Dev. Biol. 2013. 29:189–211

First published online as a Review in Advance on April 29, 2013

The *Annual Review of Cell and Developmental Biology* is online at <http://cellbio.annualreviews.org>

This article's doi:
10.1146/annurev-cellbio-101512-122324

Copyright © 2013 by Annual Reviews.
All rights reserved

Keywords

genomic imprinting, small-interfering RNA, siRNA, genome defense, transposable elements, parental conflict, DNA methylation, Polycomb group proteins

Abstract

Seeds are complex structures that unite diploid maternal tissues with filial tissues that may be haploid (gametophyte), diploid (embryo), or triploid (endosperm). Maternal tissues are predicted to favor smaller seeds than are favored by filial tissues, and filial genes of maternal origin are predicted to favor smaller seeds than are favored by filial genes of paternal origin. Consistent with these predictions, seed size is determined by an interplay between growth of maternal integuments, which limits seed size, and of filial endosperm, which promotes larger seeds. Within endosperm, genes of paternal origin favor delayed cellularization of endosperm and larger seeds, whereas genes of maternal origin favor early cellularization and smaller seeds. The ratio of maternal and paternal gene products in endosperm contributes to the failure of crosses between different ploidy levels of the same species and crosses between species. Maternally expressed small-interfering RNAs (siRNAs) are predicted to associate with growth-enhancing genes.

Contents

INTRODUCTION.....	190
CONTROL OF SEED SIZE.....	191
GENOMIC IMPRINTING IN ENDOSPERM.....	192
<i>cis</i> , <i>trans</i> , and <i>cum</i> Effects.....	193
DNA Methylation.....	195
Polycomb Group Proteins.....	195
Other Imprinted Loci.....	196
GENOMIC IMPRINTING IN EMBRYOS.....	197
INTERPLOIDY CROSSES.....	198
INTERSPECIFIC CROSSES.....	199
CONFLICTS AND PSEUDOCONFLICTS.....	200
FUNCTIONS OF siRNAS.....	201
Small-Interfering RNAs and Transposable Elements.....	201
siRNAs and Kin Conflict.....	202
Evolution of Imprinted siRNAs.....	203
ORIGIN AND SIGNIFICANCE OF ENDOSPERM.....	204

INTRODUCTION

The seed will have to be viewed not as a totally harmonious unit, but as an interdependent but sometimes contentious assemblage of relatives.

—Queller (1983)

A common mode of thinking sees parents as willing to make any sacrifice for offspring because offspring are parents' chance at genetic posterity. But this rosy picture ignores an inconvenient truth. Natural selection maximizes a parent's number of surviving offspring, not the survival of individual offspring. Parents face an evolutionary trade-off between investing less in each of a greater number of offspring or more in each of fewer offspring (Smith & Fretwell 1974).

Two ecotypes of *Arabidopsis* illustrate the trade-off between offspring size and number: *Cvi* produces 30% fewer seeds than does *Ler*, but *Cvi*'s seeds are 80% heavier (Alonso-Blanco et al. 1999). Therefore, total seed weight per plant is more similar between ecotypes than is either seed size or seed number. Similarly, overexpression of *KLU* increases *Arabidopsis* seed size without increasing total seed weight, because seeds per fruit and per plant are correspondingly reduced (Adamski et al. 2009). Thus, the size-versus-number trade-off may be expressed within individual fruits.

Mothers are selected to distribute resources among offspring to maximize the number of surviving offspring, but offspring are selected to favor themselves over sibs. Therefore, offspring will attempt to acquire more resources from mothers than mothers have been selected to supply. This phenotypic conflict will typically be resolved with the production of larger but fewer offspring than would maximize maternal fitness (Godfray 1995, Trivers 1974).

A brief flurry of activity in the 1980s applied concepts of parent-offspring conflict to problems in seed development. Different tissues within ovules (**Figure 1**) were shown to favor different trade-offs between seed size and seed number. Maternal tissues were predicted to abort a subset of seeds that contained embryos of low quality, or were surplus to the number mothers could provision, and

Ecotype: the equivalent of an inbred line, propagated by self-fertilization

Ovule: the maternal multicellular structure that develops into a seed after fertilization

Embryo: product of the fertilization of the egg nucleus of the female gametophyte by one of the two sperm released by a pollen tube




Figure 1

(a) Before fertilization, the diploid ovule contains a haploid female gametophyte. A male gametophyte (pollen tube) grows to the ovule and releases two sperm. One sperm nucleus fuses with the egg nucleus to produce a zygote. The other sperm fuses with two haploid nuclei of the central cell to produce a triploid primary endosperm nucleus. (b) After fertilization, ovules are known as seeds. A seed consists of a diploid seed coat derived from the maternal integuments, a diploid embryo derived from the zygote, and a triploid endosperm derived from the primary endosperm nucleus. In the figure, a tissue containing x maternal genomes and y paternal genomes is labeled as $xm:yp$. The maternal sporophyte and seed coat are labeled with asterisks ($1m^*:1p^*$) to indicate that these are maternal and paternal genomes of the generation preceding the embryo and endosperm.

to constrain the growth of seeds that were provisioned. Conversely, filial tissues were predicted to express adaptations to evade abortion and promote seed growth (Haig & Westoby 1988; Queller 1983, 1989; Westoby & Rice 1982).

The unusual genetic makeup of endosperm, triploid with two maternal genomes and one paternal genome ($2m:1p$), was a particular focus of attention. When endosperm was compared with an embryo, its extra maternal genome was interpreted as reducing filial demands and conferring greater maternal control (Westoby & Rice 1982), whereas, when endosperm was compared with a female gametophyte, its extra paternal genome was interpreted as favoring more aggressive acquisition of resources (Queller 1983). The parental-conflict theory of the evolution of genomic imprinting emerged from consideration of this contrast (Haig & Westoby 1989).

CONTROL OF SEED SIZE

Genetic data on seed size come mostly from *Arabidopsis* and the major grain cereals. These taxa may be atypical, because parent-offspring conflict should be attenuated in predominantly self-fertilizing *Arabidopsis thaliana* (De Jong et al. 2011) and because grain cereals have been subject to intense artificial selection to increase seed yields.

Arabidopsis seed volume is determined early, during rapid expansion of the central vacuole of syncytial endosperm. After endosperm cellularizes, the embryo expands, crushing the endosperm and accumulating food reserves, but there is little further increase in seed volume. Thus, seed

Endosperm: product of the fertilization of the central cell of the female gametophyte by one of the two sperm released by a pollen tube; the central cell usually contains two haploid nuclei, both of which fuse with the sperm nucleus to form the triploid primary endosperm nucleus

Female gametophyte: the multicellular haploid plant enclosed within an ovule; fertilization of its egg cell gives rise to an embryo, and fertilization of its central cell gives rise to endosperm

Invertase: enzyme that catalyzes hydrolysis of sucrose to fructose and glucose

Integument: a maternal diploid tissue surrounding an ovule; integuments give rise to the seed coat, and most angiosperms possess an inner and outer integument

Transfer cells: cells with elaborate wall ingrowths that dramatically increase the cell membrane's surface area; transfer cells are generally believed to play a role in secretion and/or absorption

Sporophyte: the diploid plant that produces haploid spores by meiosis

maturation is associated with a rapid early increase in fresh weight, accompanied by a slower, but steady, increase in dry weight (Baud et al. 2002, Brown et al. 1999, Mansfield & Briarty 1992). Seed size in grasses is also determined by early endosperm expansion followed by deposition of food reserves, but, unlike in *Arabidopsis*, embryos remain relatively small, and reserves are deposited in persistent endosperm (Olsen 2004).

During seed expansion of *Brassica napus*, invertase splits sucrose into hexose sugars, doubling osmotic strength. The osmotically driven influx of water is conjectured to facilitate rapid expansion of the central endosperm vacuole (Morley-Smith et al. 2008). In mechanical terms, endosperm turgor exerts tension on integuments, with the stiffness of integument cell walls determining the compression exerted on endosperm. Seed volume would then reflect, in part, how integuments respond to tension and endosperm to compression. *Mn1* encodes the invertase expressed in basal endosperm transfer cells of maize during endosperm expansion (Kang et al. 2010). Loss of *Mn1* activity leads to endosperms with fewer and smaller cells (Vilhar et al. 2002). The causes of growth retardation are unknown, but reduced turgor may contribute.

Endosperm expansion and integument growth interact to determine seed volume. *Arabidopsis* mutations with maternal effects on seed size commonly affect the proliferation or elongation of integument cells (Adamski et al. 2009, FitzGerald et al. 2008, Garcia et al. 2005, Hughes et al. 2008, Li et al. 2008, Ohto et al. 2009, Schruff et al. 2006), whereas mutations with filial effects on seed size commonly affect the timing of endosperm cellularization (Garcia et al. 2003, Kang et al. 2008, Luo et al. 2005, Wang et al. 2010, Zhou et al. 2009).

Endosperm and integument growth are coupled. Maternal *ttg2* mutations cause reduced elongation of integuments and, as a filial response, precocious cellularization of endosperm, whereas filial *baiku* mutations cause reduced proliferation of endosperm and, as a maternal response, reduced elongation of integuments (Berger et al. 2006, Garcia et al. 2005). The emerging picture, of maternal constraints on filial growth, is broadly consistent with kin-conflict theory (Queller 1983, Westoby & Rice 1982).

GENOMIC IMPRINTING IN ENDOSPERM

The genetic basis of parent-offspring conflict is subtle. If a sporophyte is heterozygous for alleles at a locus expressed in integuments, then all ovules will exhibit the same pattern of expression, independent of which allele is inherited by the enclosed filial tissues. However, if the locus is expressed in filial tissues (female gametophytes, embryos, or endosperms), then the effects of maternal alleles will segregate among ovules depending on which allele a particular tissue inherits. Therefore, maternal alleles expressed in sporophytic tissues are selected to maximize the aggregate fitness of all embryos, whereas maternal alleles expressed in filial tissues are selected to favor embryos with their copies over embryos without. A similar argument applies to paternal alleles expressed in filial tissues, but with a twist. Filial tissues of different seeds on a maternal sporophyte are less likely to share genes of paternal origin (patrigenes) than genes of maternal origin (matrigenes) when mothers produce offspring with multiple fathers. Therefore, patrigenes favor larger seeds than do matrigenes because competition among patrigenes is less constrained by costs to sibs (Haig & Westoby 1989; the matrigenes/patrigenes terminology was proposed by Queller 2003).

Maternal genes (expressed in maternal sporophytes) and matrigenes (expressed in filial tissues) are subject to different selective forces because alleles segregate at meiosis. Paternal genes (expressed in fathers) and patrigenes (expressed in filial tissues) similarly favor different outcomes. Parental conflict is a convenient label, but it is a misnomer if interpreted to mean that conflict between matrigenes and patrigenes is the same as conflict between maternal and paternal genes.

Matrigenes descend from an allele present in a megaspore and benefit from the same outcomes, whether expressed in female gametophytes before fertilization or from maternally derived copies in embryos or endosperms after fertilization. Patrigenes descend from an allele present in a microspore and benefit from the same outcomes, whether expressed in male gametophytes or from paternally derived copies in embryos or endosperms. A gametophytic effect could be explained by persistence of transcripts from before fertilization or by parent-specific expression after fertilization (Curtis & Grossniklaus 2008). This makes a big difference if one's interest is mechanism, but it makes no difference if one's interest is function. It is the same gene with the same interests, whether it is transcribed in female gametophytes before fertilization or from maternally derived copies after fertilization. Kin-conflict theory distinguishes between gene expression before and after meiosis but not between gametophytic expression before fertilization and imprinted expression after fertilization.

Whether matrigenes and patrigenes pursue divergent goals in embryos and endosperms depends on genomic imprinting. Imprinted expression allows a gene to adopt different matrigenic and patrigenic roles, whereas unimprinted expression forces an evolutionary compromise. Kin-conflict theory predicts that maternally expressed genes (MEGs) should reduce filial demands on mothers, whereas paternally expressed genes (PEGs) should increase filial demands, with the effects of unimprinted genes falling somewhere in the middle (Haig & Westoby 1989, 1991). In a completely self-fertilizing plant, matrigenes, patrigenes, and parental genes favor the same seed size.

Megaspore:

a product of female meiosis; megaspores produce female gametophytes by mitosis

Microspore:

a product of male meiosis; microspores produce male gametophytes (pollen) by mitosis

Function: “function of X” is used in this review as shorthand for “reason why natural selection favors and maintains X”

***cis*, *trans*, and *cum* Effects**

Gene expression is commonly viewed as jointly determined by *cis* effects, intrinsic to an expressed sequence's haplotype, and *trans* effects, exerted by the products of other genes (Figure 2). Epigenetic modifications in *cis* and in *trans* are subject to different selective forces when meiosis intervenes between the establishment of a heritable modification and its effect on gene expression. Suppose a parent is heterozygous for alleles at a locus that modifies imprints in *cis*. An imprint's effect on gene expression will be experienced only by offspring who inherit the allele responsible for the imprint. Therefore, epigenetic effects in *cis* will evolve to favor offspring who inherit the allele responsible for an effect at the expense of siblings who inherit alternative alleles (Figure 3a).




Figure 2

Expression of *A* is determined by a mixture of *cis*, *trans*, and *cum* effects. *cis* effects are mediated by linked sites on the same chromosome and therefore affect only the linked copy of *A*. *trans* effects are mediated by gene products (proteins or noncoding RNAs) and affect copies of *A* on all homologous chromosomes. A *cum* effect is an effect of *A* on its own copies on other chromosomes.




Figure 3

(a) A modifier (*purple*) causes an epigenetic modification (*red*) in *cis*. The modifier and modification are inherited by the same set of offspring. (b) A modifier (*purple*) causes an epigenetic modification (*red*) in *trans*. Offspring with and without the modifier inherit the modification. (c) A modifier (*purple*) causes an allele-specific epigenetic modification (*red*) in *trans*. The modifier and modification segregate independently to offspring.

Suppose, instead, that the parent is heterozygous at a *trans*-acting locus that modifies imprints on both alleles at a second locus that is expressed in offspring. All of the parent's offspring will exhibit the same pattern of expression, independent of which allele at the *trans*-acting locus a particular offspring inherits (**Figure 3b**). If, however, alleles at the imprinted locus are differentially modified by the *trans*-acting locus, then these *cis* effects will segregate among offspring independently of alleles at the *trans*-acting locus (**Figure 3c**; assuming the loci are unlinked). In either case, *trans*-acting loci will evolve to favor parental interests.

Mammals establish imprints through *trans*-acting genes expressed in parental germ cells before meiosis. Therefore, conflict is possible between imprinting genes expressed in parents and imprinted genes expressed in offspring, with imprinted loci selected to reinterpret in *cis* parental instructions received in *trans*. Conflict similarly exists between imprinted and unimprinted loci expressed in offspring (Burt & Trivers 1998). For these reasons, natural selection at imprinted loci is expected to favor *cis* control. This may explain why imprinted loci in mammals often occur in *cis*-regulated clusters (Verona et al. 2003).

Angiosperms establish imprints in haploid gametophytes after meiosis. Therefore, *trans* effects of an imprinting locus are experienced by single alleles at imprinted loci. Conflict is absent because imprinter and imprintee are inherited by the same set of offspring. Therefore, natural selection does not favor *cis* over *trans* control at imprinted loci. This may explain why mammal-like clusters

of imprinted loci have not been found in plants (Luo et al. 2011, Wolff et al. 2011; but see Zhang et al. 2011).

A third set of effects are subject to different selective forces from *cis* and *trans* effects. I will call these *cum* effects, for want of a better Latin preposition. A *cum* effect, or “*trans*-homolog interaction” (Hollick 2012), is an effect of a sequence on the expression of another copy of the same sequence (**Figure 2**). Genes with *cum* effects can respond to their own dosage and detect whether their copies are inherited from the other parent, which creates opportunities for collusion between maternally and paternally inherited alleles. Processes by which genes recognize their own copies have been called green beard effects, after a genetically encoded tag (the eponymous green beard) that allows altruists to recognize and cooperate with each other (Haig 1996, West & Gardner 2010). Newly arisen *cum* effects can function as intragenomic green beards, although the reliability of self-recognition decays as mutation creates variants that retain the tag but are no longer self. Existing models of kin conflict during seed development have not considered *cum* effects.

DNA Methylation

Maternally derived genomes of *Arabidopsis* endosperm are extensively demethylated relative to the paternally derived genome of endosperm and the genomes of embryos and vegetative cells (Gehring et al. 2009, Hsieh et al. 2009). Differential methylation is established by coordinated expression of *DME*, a gene that encodes a DNA demethylase, and *MET1*, a gene that encodes a DNA methyltransferase. *DME* is expressed in central cells but in neither sperm nor endosperm (Choi et al. 2002, Gehring et al. 2006, Schoft et al. 2011). *MET1* is expressed in sperm but not in central cells (Jullien et al. 2012, Xiao et al. 2006). Therefore, at the time of fertilization, central cells are relatively demethylated, whereas sperm are relatively methylated, and this difference between maternal and paternal genomes is maintained after fertilization (Köhler et al. 2012). The methylated paternal genome and demethylated maternal genomes of *Arabidopsis* endosperm contrast with the methylated maternal genome and demethylated paternal genome of mouse zygotes (Wu & Zhang 2012).

DME is expressed in central cells but not in sperm cells, whereas *MET1* is expressed in sperm cells but not in central cells. Therefore, kin-conflict theory predicts *DME* should reduce endosperm proliferation, whereas *MET1* should promote endosperm growth. Consistent with these predictions, seeds with maternal *dme* mutations have enlarged endosperm (Choi et al. 2002), whereas seeds with paternal *met1* mutations produce small, precociously cellularized endosperm (Xiao et al. 2006). A simple story in which *DME* and *MET1* combine to establish different epigenetic states of maternal and paternal chromosomes cannot explain all imprinting in *Arabidopsis* endosperm, because imprinted expression of *MEA* and a large class of small-interfering RNAs (siRNAs) is maintained in the presence of maternal *dme* and paternal *met1* mutations (Mosher et al. 2011, Wöhrmann et al. 2012). Rice endosperm does not undergo global demethylation, nor have orthologs of *DME* been detected in the genomes of rice or other monocots (Zemach et al. 2010). Differentially methylated regions of maize endosperm are maternally hypomethylated and paternally methylated, but differential methylation is more localized than in *Arabidopsis* (Lauria et al. 2004).

Polycomb Group Proteins

Polycomb repressive complex 2 (PRC2) catalyzes trimethylation of lysine 27 residues on histone H3. H3K27me3 is frequently associated with transcriptional repression of associated DNA. Three

subunits of PRC2 (*MEA*, *FIS2*, and *FIE*) exhibit matrigenic expression in *Arabidopsis* endosperm (Rodrigues et al. 2010).

MEA is expressed in female gametophytes and from maternally derived alleles in postfertilization endosperm (Vielle-Calzada et al. 1999). A 200-bp sequence is necessary and sufficient for imprinted expression of *MEA* (Wöhrmann et al. 2012). *MEA* protein binds directly to *MEA* promoters and reduces transcription of maternal *MEA* alleles in endosperm. As a consequence of the absence of functional *MEA*, *mea* mutations are associated with dramatic upregulation of their own mRNA (Baroux et al. 2006). *MEA* also acts as part of PRC2 to repress paternal *MEA* transcription in endosperm (Gehring et al. 2006, Jullien et al. 2006).

FIS2 is expressed in central cells before fertilization and from maternally derived alleles in endosperm after fertilization (Luo et al. 2000). Therefore, *FIS2* and *MEA* have similar patterns of expression, but, unlike *MEA*, imprinting of *FIS2* depends on DNA methylation (Jullien et al. 2006). *FIE* is expressed exclusively from maternal alleles in early endosperm but from maternal and paternal alleles in later endosperm (Luo et al. 2000, Yadegari et al. 2000). Despite this phase of biallelic expression, patrigenic *FIE* does not rescue the endosperm overproliferation and seed abortion caused by matrigenic *fie* mutations (Vinkenoog et al. 2000). *fie* mutations are associated with reduced accumulation of their own transcripts, in contrast to the effects of *mea* mutations that increase *mea* transcripts (Baroux et al. 2006). Patrigenic *fie* mutations cause patrigenic expression of *MEA* in endosperm (Jullien et al. 2006).

Seeds with matrigenic *mea*, *fis2*, or *fie* mutations initiate endosperm development without fertilization (Chaudhury et al. 1997; Luo et al. 1999, 2000) and exhibit prolonged proliferation and delayed cellularization of endosperm if central cells are fertilized (Grossniklaus et al. 1998, Guitton & Berger 2011, Kiyosue et al. 1999, Köhler & Makarevich 2006). Thus, wild-type PRC2 prevents endosperm proliferation before fertilization and accelerates differentiation of endosperm after fertilization (Ingouff et al. 2005b). These functions are consistent with predictions that MEGs will be associated with restraints on endosperm growth.

PRC2 genes of *Arabidopsis* and grasses lack one-to-one correspondence because of a complex history of gene duplication. The genomes of rice and maize lack obvious orthologs of *MEA* and *FIS2* (Chen et al. 2009, Spillane et al. 2007) but possess two *FIE*-related genes, at least one of which is maternally expressed in endosperm (Danilevskaya et al. 2003, Dickinson et al. 2012, Luo et al. 2009, Springer et al. 2002). Unlike *fie* mutations in *Arabidopsis*, disruption of *FIE1* in rice does not cause endosperm to develop without fertilization (Luo et al. 2009). Thus, the expression and function of imprinted genes are not conserved.

Three papers on molecular evolution of *MEA* collectively found stronger selection for new variants in outcrossing *A. lyrata* than in self-fertilizing *A. thaliana*, with evidence of balancing selection at the *MEA* promoter of *A. lyrata* (Kawabe et al. 2007, Miyake et al. 2009, Spillane et al. 2007). McVean & Hurst (1997) claimed the kin-conflict hypothesis entails an evolutionary arms race that should leave a signature of accelerated evolution in the sequence of imprinted genes. However, an explicit model of the joint evolution of oppositely imprinted loci with antagonistic effects found a stable equilibrium at which there was no further change in expression levels (Wilkins & Haig 2001). Red Queen dynamics (running as fast as you can to stay in place) are not an inevitable outcome of kin conflict.

Other Imprinted Loci

Deep sequencing of mRNA from reciprocal crosses between divergent strains of *Arabidopsis* (Gehring et al. 2011, Hsieh et al. 2011, Wolff et al. 2011), rice (Luo et al. 2011), and maize (Waters et al. 2011, Zhang et al. 2011) has identified numerous loci preferentially expressed from

maternal or paternal alleles during seed development. These screens have substantially increased the number of candidate imprinted loci. Lists from different screens, even within the same species, show minimal overlap (Pignatta & Gehring 2012).

Little is known about the phenotypic effects of most imprinted genes. *FWA* and *AGL36*, for example, are MEGs expressed in endosperm, but their loss of function lacks an obvious phenotype in endosperm (Kinoshita et al. 2004, Lippman et al. 2004, Shirzadi et al. 2011). Therefore, whether these genes' functions support the kin-conflict hypothesis cannot be evaluated. Three imprinted loci with endosperm phenotypes are considered below. *PHE1* (*Arabidopsis*) is a PEG that exhibits a phenotype consistent with the kin-conflict hypothesis. *FH5* (*Arabidopsis*) and *Meg1* (maize) are MEGs whose phenotypes have been considered to be inconsistent with the hypothesis.

PHE1 is expressed from paternal alleles in syncytial endosperm. Maternal alleles are normally repressed by PRC2 but are reactivated in *mea* endosperms (Köhler et al. 2005; Makarevich et al. 2006, 2008; Villar et al. 2009). *mea* seeds abort after prolonged nuclear proliferation of endosperm, but this phenotype is partially rescued by reduced expression of *PHE1*. The rescued seeds are larger than normal (Köhler et al. 2003). These data are compatible with a model in which paternally expressed *PHE1* promotes nuclear proliferation in endosperm and larger seed size, but its expression is attenuated by maternally expressed *MEA* (Schubert & Goodrich 2003). Another MEG (*At1g59930*) encodes a truncated PHE1-related protein that may act as a PHE1 antagonist (Hsieh et al. 2011).

FH5 is expressed from maternal alleles in chalazal endosperm (FitzGerald et al. 2009) with cellularization delayed in *fb5* seeds (Ingouff et al. 2005a). This phenotype has been interpreted as contradicting the kin-conflict hypothesis (Jullien & Berger 2010) but appears consistent with a model in which conflict is mediated by antagonism between endosperm proliferation (promoted by PEGs such as *PHE1*) and differentiation (promoted by MEGs such as *FH5*).

Meg1 is expressed from maternal alleles in just-cellularized endosperm but from both alleles at later stages of development (Gutiérrez-Marcos et al. 2004). Its expression is necessary and sufficient for formation of endosperm transfer cells. Knockdown of *Meg1* is associated with reduced seed weight, whereas increased *Meg1* is associated with a dosage-sensitive increase in seed weight (Costa et al. 2012). Therefore, *Meg1* has been considered to contradict the parental-conflict hypothesis (Ikeda 2012, Jiang & Köhler 2012, Li & Berger 2012).

Effects of *Meg1* during the early phase of imprinted expression and the later phase of biallelic expression are entangled in Costa et al.'s (2012) study. Significantly, imprinted *Meg1* transgenes increase seed weight when inherited from fathers. Thus, at least some of *Meg1*'s effects on seed weight result from its biallelic expression, but the parental-conflict hypothesis addresses only effects of imprinted expression. Differentiation of transfer cells is determined early in syncytial development (Costa et al. 2003). Therefore, the hypothesis must address this aspect of phenotype. Imprinted expression of *Meg1* could be viewed as another expression of tension between MEG-promoted differentiation and PEG-promoted proliferation. This interpretation remains to be tested.

GENOMIC IMPRINTING IN EMBRYOS

Kin-conflict theory predicts imprinting of genes in endosperm but is equivocal about imprinting of genes in embryos. Seed volume is determined by maternal-endosperm interactions before food reserves are deposited, but the metabolic load on mothers, and competition among sibs, may be influenced by sink strengths during the filling phase. Whether embryos are passive observers or active participants in conflict over seed weight will depend on details of seed development and physiology.

A screen for imprinted loci in rice identified 262 candidates in endosperm but only three in embryos (Luo et al. 2011). Similar studies in maize (Waters et al. 2011) and *Arabidopsis* (Gehring et al. 2011) detected few imprinted genes in embryos. One gene (*mee1*) has been shown to be maternally expressed in early embryos and endosperm of maize (Jahnke & Scholten 2009).

The paternal genome of *Arabidopsis* embryos is variously reported to be inactive during early embryogenesis (Autran et al. 2011) or to be active in two-celled embryos (Nodine & Bartel 2012). The latter report found little carryover of mRNA from gametophytes in early embryos but detected 77 transcripts with significant matrigenic bias and 44 transcripts with significant patrigenic bias, which suggests short-lived imprinting effects.

SSP mRNAs carried by *Arabidopsis* sperm are translated in zygotes and regulate the first asymmetric cell division that establishes apical (embryo) and basal (suspensor) cell fates (Bayer et al. 2009). SSP acquired its current function sometime after a whole-genome duplication in an ancestor of the Brassicaceae (Liu & Adams 2010). mRNAs transcribed in male gametophytes and delivered to zygotes via sperm are subject to the same selective forces as PEGs transcribed in embryos after fertilization.

INTERPLOIDY CROSSES

Failure of crosses between diploids ($2x$) and their own autotetraploids ($4x$) posed a puzzle for plant breeders. Viable seed was produced when diploids were crossed with diploids (embryo $2x$, endosperm $3x$) or tetraploids with tetraploids (embryo $4x$, endosperm $6x$), but endosperm development was grossly perturbed when diploids were crossed with tetraploids. Embryos from interploidy crosses were triploid ($3x$), intermediate between viable $2x$ and $4x$ embryos, and endosperms were $4x$ or $5x$, intermediate between viable $3x$ and $6x$ endosperms. So why did the crosses fail? Various hypotheses to explain triploid block invoked requirements for particular ploidy ratios of seed coat, embryo, and endosperm, but these hypotheses were ad hoc attempts to fit the data without theoretical reasons why ploidy ratios should make a difference (Haig & Westoby 1991).

An elegant study in maize solved this puzzle by showing that normal endosperm development requires two maternal genomes for each paternal genome. Central cells of *ig* female gametophytes contain variable numbers of supernumerary nuclei. Fertilization of *ig* central cells by $1x$ or $2x$ pollen creates endosperms with a range of maternal and paternal ploidies. Most combinations abort. Normal endosperms are either $3x$ with two maternal and one paternal genome ($2m:1p$) or $6x$ ($4m:2p$). Significantly, $6x$ endosperms abort ($5m:1p$) or are well formed ($4m:2p$), depending on the parental-genome ratio (Lin 1984).

Haig & Westoby (1991) proposed that seed phenotypes in crosses between $2x$ and $4x$ parents were explicable in terms of antagonistic actions of maternal and paternal genomes, with MEGs restraining and PEGs promoting endosperm growth. When mothers are $4x$ and fathers $2x$, endosperms are $5x$ ($4m:1p$) and seeds are typically small with prematurely cellularized endosperm. The authors called this maternal excess. In the reciprocal cross, endosperms are $4x$ and seeds are often full-sized but shriveled because endosperm fails to cellularize. The authors called this paternal excess.

Viable seeds are produced in interploidy crosses involving *Arabidopsis* accessions *Ler* and C24 (Scott et al. 1998). Seeds with $4m:1p$ endosperms are lighter than normal, with slow mitosis in endosperm, whereas seeds with $2m:2p$ endosperms are heavier than normal, with accelerated endosperm mitosis. A plausible explanation for why $2m:2p$ endosperms are viable in *A. thaliana*, but nonviable in outcrossing relatives of *Arabidopsis* (Stoute et al. 2012) and most other species, is that conflict between matrigenes and patrigenes is attenuated in *A. thaliana* because of high rates of self-fertilization (Scott et al. 1998). In this view, imbalance in $2m:2p$ endosperms of

outcrossing species causes seed abortion, but the underlying growth enhancement of paternal excess is uncovered in self-fertilizing *Arabidopsis*, where imbalance is less severe.

Maternal sporophytic factors contribute to seed abortion in interploidy crosses. Fewer plump seeds are produced in interploidy crosses by using 4x Col rather than 4x *Ler* as father (paternal effect) and 2x Col rather than 2x *Ler* as mother (maternal effect). Recombinant inbred lines between Col and *Ler* mapped the maternal effect to a quantitative trait locus containing *TTG2*. Seed lethality in crosses with 4x Col fathers was rescued if 2x mothers were *ttg2* homozygotes (Dilkes et al. 2008).

Most seeds abort in interploidy crosses in maize; 2m:2p endosperms show prolonged proliferation with delayed onset of endoreduplication and fail to form a transfer cell layer (Charlton et al. 1995, Leblanc et al. 2002, Pennington et al. 2008), and 4m:1p endosperms show reduced mitotic proliferation with premature onset of endoreduplication (Leblanc et al. 2002, Pennington et al. 2008). Maternal excess causes early silencing of genes associated with endosperm proliferation and early activation of genes associated with grain filling. Paternal excess has opposite effects (Li & Dickinson 2010).

These studies suggest that maternal genomes of endosperm promote early differentiation with foreshortened nuclear proliferation and that paternal genomes promote delayed differentiation with prolonged nuclear proliferation. Thus, the marginal effect of matrigenic expression would be to reduce seed size, and the marginal effect of patrigenic expression would be to increase seed size. These effects are sometimes obscured by abortion of seeds with strongly perturbed development. Factors expressed in sporophytic tissues, with balanced maternal-to-paternal ratios, contribute to failure of seeds with imbalanced ratios in filial tissues.

INTERSPECIFIC CROSSES

Failure of crosses between species sometimes resembles failure of crosses between different ploidy levels within species, with features of maternal excess observed in one direction of a cross and features of paternal excess in the reciprocal cross (Bushell et al. 2003, Gutiérrez-Marcos et al. 2003, Johnston et al. 1980, Köhler & Kradolfer 2011). Success or failure of crosses between *Avena* species could be predicted by assigning each parent an activating value (AV) for its sperm nuclei and an equal response value (RV) for each nucleus of the central cell (Nishiyama & Yabuno 1978, 1979). Species with similar AVs crossed readily, but development was perturbed in crosses between species with markedly different AVs. Normal endosperm development required an activation index (AI) = AV/2RV in the range of 0.3 to 0.8; seeds were small and unviable if the AI was less than 0.2 and small but viable for AIs between 0.2 and 0.3, but they were shriveled and empty for AIs greater than 0.8.

A similar scheme was proposed to explain crosses between *Solanum* species. Each species was assigned an endosperm balance number (EBN) determined by its behavior in crosses to a reference species. Normal endosperm required a 2:1 ratio of maternal to paternal EBNs (Johnston et al. 1980). Failed crosses could sometimes be rescued by changing the ploidy of one of the parents to bring the ratio closer to 2:1 (Johnston & Hanneman 1982).

Haig & Westoby (1991) proposed that divergent expression of imprinted genes between species can result in developmental incompatibilities in hybrid endosperm, because the balance of maternal and paternal products is perturbed. If divergence at multiple loci were haphazard, then increasing the ploidy of one parent might correct the ratio of products for some loci but exacerbate the problem at other loci. Therefore, we were puzzled why crossing behavior could be summarized by a single number (EBN or AI). Such indices suggested the effects of multiple imprinted loci aligned on a single developmental axis.

Mating system provides a possible solution to this puzzle. Conflict between matrigenes and patrigenes becomes less intense as single paternity of a mother's offspring becomes more common (Kondoh & Higashi 2000). Therefore, the divergence in mating system could explain why multiple imprinted loci exhibit coordinated effects in hybrid seeds. A literature review of crosses between outcrossing and self-fertilizing plants found features of paternal excess when outcrossers were fathers but of maternal excess when outcrossers were mothers (Brandvain & Haig 2005).

Endosperms exhibit severe paternal excess when *A. thaliana* is crossed as seed parent to *A. arenosa*, but the cross succeeds when 4x *A. thaliana* is substituted as seed parent. In reciprocal backcrosses of the latter hybrids to 4x *A. thaliana*, genomes of *A. arenosa* promoted early endosperm cellularization when contributed maternally but delayed endosperm cellularization when contributed paternally. Therefore, outcrossing *A. arenosa* was inferred to express greater genomic strength than self-fertilizing *A. thaliana* (Bushell et al. 2003), consistent with expectations from the divergence in mating system. Studies of hybrid *Arabidopsis* endosperm have shown extensive disruption of gene expression, including loss of imprinting of *PHE1* and *MEA* (Josefsson et al. 2006, Walia et al. 2009).

CONFLICTS AND PSEUDOCONFLICTS

Two kinds of answer can be given to the question of why a flower is red. The first explains redness in terms of mechanism: Genes of the anthocyanin pathway are activated in petals. The second explains redness in terms of function: Red flowers attract hummingbirds that move pollen from flower to flower. Different senses of why are addressed by the two answers, and few would argue that the truth of one negates that of the other. Unfortunately, explanations of mechanism and function are frequently confused when discussing the why of genomic imprinting.

The maternal-offspring-coadaptation hypothesis explains the function of imprinted expression in terms of coordination rather than conflict (Curley et al. 2004, Keverne & Curley 2008, Swaney et al. 2007). The hypothesis is supported by a model of an epistatic interaction between an unimprinted gene in mothers and its imprinted or unimprinted copies in offspring (transgenerational *cum* effect). In this model, homozygous mothers benefit from inactivation of the paternal allele of their heterozygous offspring (Wolf & Hager 2006). At the time of writing, none of the imprinted genes that are claimed to support the hypothesis have been shown to conform to the model's rather specific assumptions (Haig 2013).

The differential-dosage hypothesis proposes that many mechanisms, not just imprinted expression, cause parental effects in seed development (Dilkes & Comai 2004). Undoubtedly, parental effects exist that do not involve imprinted genes, but the kin-conflict hypothesis neither denies their existence nor purports to explain them, whereas the differential-dosage hypothesis strives to understand mechanisms of seed development but does not purport to explain why parent-specific expression evolved. Presentation of these hypotheses as competitors confuses a hypothesis about mechanisms with one about functions.

The genome-defense hypothesis proposes that DNA methylation and RNA interference have evolved to control the spread of genomic parasites and that these mechanisms are the means whereby genes acquire imprinted expression. The hypothesis has two separable components. The first posits that inactivation of selfish genetic elements is an important function, perhaps the primary function, of epigenetic gene silencing (Köhler & Weinhofer-Molisch 2010, Matzke et al. 1996). The second posits that some genes have acquired imprinted expression because of their proximity, or resemblance, to sequences subject to processes of genome defense (Gehring et al. 2009, Slotkin & Martienssen 2007). Thus, a hypothesis about the function of epigenetic mechanisms is combined with a hypothesis about the mechanism of imprinted expression.

Neither hypothesis directly challenges the kin-conflict hypothesis, which addresses the function of imprinted expression.

Mechanisms of gene silencing may have evolved for reasons of defense, but they have been employed by other genes to achieve parent-specific expression because of conflict (Gehring et al. 2009, Köhler & Weinhofer-Molisch 2010). Male and female germ lines will possess different vulnerabilities to transposable elements (TEs) and different repertoires of host defense. Sex-specific adaptations of TEs, sex-specific counteradaptations of hosts, and collateral effects of these adaptations provide a source of parent-specific variation in gene expression on which natural selection can act to mediate kin conflict (Haig 2012).

Explanations of mechanism do not obviate the need for explanations of function (why we observe this mechanism and not others) or lack of function. If the insertion of a transposable element confers imprinted expression upon an allele in *cis*, then the initially rare imprinted allele becomes established in the population in one of two ways. Either imprinted expression confers a benefit that causes the allele to sweep to high frequency under natural selection or imprinted expression has minimal effects on fitness and the allele drifts to high frequency through random processes. In the first case, the reasons why an imprinted allele is favored and maintained by natural selection relative to unimprinted alleles comprise the function of imprinted expression. In the second case, imprinted expression does not have a function but is a side effect of defense mechanisms (Haig & Trivers 1995).

What is the evolutionary history of genomic imprinting? What are the mechanisms of imprinting, and what are the functions of these mechanisms? Why have some sequences, but not others, evolved to use these mechanisms to achieve imprinted expression? These are all important questions. Much needless argument would be avoided by paying close attention to the questions addressed by each hypothesis. Conflict and coadaptation hypotheses both purport to explain the function of imprinted gene expression. Differential dosage and genome defense are sometimes presented as rivals of conflict but, for the most part, address different questions.

FUNCTIONS OF siRNAS

Small-Interfering RNAs and Transposable Elements

siRNAs target DNA methylation and histone modification to complementary DNA sequences and, by this means, exert *trans* or *cum* effects on gene expression. siRNAs complementary to TEs are believed to function in host defense (Lisch & Slotkin 2011). Most of the machinery that synthesizes siRNAs acts in *trans* and should evolve to promote host fitness and reduce TE activity, but how do the sequences recognized by this machinery evolve in *cis* and *cum*? The question of whether an siRNA transcribed from a TE's own sequence is an adaptation of the host or the parasite is not simple.

Natural selection after insertion of a TE selects for variants that enhance fitness of the haplotype on which the TE resides, but acts of insertion select for transposition-competent TEs. Therefore, the lineage of a recently inserted TE will have been subject to both forms of selection. The footprints left by a mobile TE as it wanders through the genome are either erased by selective elimination of costly insertions or remain as domesticated remnants subject to degradation of their ability to transpose (Haig 2012). siRNA-encoding sequences of domesticated TEs can be considered host adaptations if they are selectively maintained because of their ability to silence their host sequence or related TEs. siRNAs could also be considered adaptations of TEs if self-restrained TEs spread more readily than unrestrained TEs.

Plasmodesmata:
narrow cytoplasmic
connections between
neighboring cells

Self-restraint of TEs is considered unlikely to evolve in outbred plants but more likely to evolve in selfing plants (Charlesworth & Langley 1986). Insertions often reduce fitness. TEs of outcrossing plants that generate new insertions soon segregate away from their costly progeny, just as less deleterious insertions segregate away from their more costly brethren. By contrast, the recent copies of an active TE of a self-fertilizing plant are yoked together for many generations before parting company after rare outcrossing. An overly active TE would foul its own nest. For this reason, TEs may evolve to be less virulent in selfing plants.

Once a TE is homozygous because of selfing, it is transmitted to all of a sporophyte's progeny. Propagation of new insertions might be facilitated by restricting transposition to occasions on which the TE is rendered heterozygous by outcrossing. Dosage-sensitive responses to its own *cum*-acting siRNA might allow a canny TE to recognize, and act appropriately toward, its own copies on other chromosomes.

Arabidopsis may use strategic bursts of TE activity to immunize its genome against proliferation of TEs. In this scenario, TEs are activated in cells that do not contribute to future generations; siRNAs are generated against the active TEs and then exported to nuclei of the germ lineage to silence endogenous TEs. On the male side, TEs are activated in vegetative nuclei of pollen grains and 21-nucleotide (nt) siRNAs exported to sperm nuclei (Slotkin et al. 2009). On the female side, TEs are activated in endosperm and 24-nt siRNAs exported to embryos (Hsieh et al. 2009, Mosher & Melnyk 2010). The route by which siRNAs move from endosperm to embryo in the latter hypothesis is unclear, because plasmodesmata are absent between embryo and endosperm (Mansfield & Briarty 1991, Molnar et al. 2010).

siRNAs and Kin Conflict

Imbalances between 21-nt siRNAs contributed by sperm and TEs of central cells and imbalances between 24-nt siRNAs contributed by central cells and TEs of sperm have been proposed to underlie failures of interploidy and interspecific crosses (Martienssen 2010; also see Josefsson et al. 2006). These effects could also be explained by siRNAs directly targeting genes involved in endosperm growth and differentiation. siRNAs transcribed in male and female gametophytes are subject to the same selective forces as MEGs and PEGs. Because siRNAs reduce expression of the genes from which they are transcribed, matrigenically expressed siRNAs are predicted to accumulate in genes whose expression increases endosperm growth, and patrigenically expressed siRNAs are predicted to accumulate in genes whose expression inhibits endosperm growth.

A highly diverse and abundant class of 24-nt siRNAs are expressed in *Arabidopsis* female gametophytes before fertilization and from maternal chromosomes of endosperm after fertilization (Mosher et al. 2009). Many of these siRNAs are not associated with TEs, and mutations that abolish their expression do not reactivate TEs (Mosher 2010, Mosher & Melnyk 2010). In reciprocal crosses between $2x$ and $4x$ *Arabidopsis*, 24-nt siRNAs are increased in maternal-excess endosperms but decreased in paternal-excess endosperms, and they appear to promote precocious cellularization by targeting genes that promote proliferation (Lu et al. 2012). These observations are consistent with kin-conflict predictions. However, maternal mutations that abolish expression of 24-nt siRNAs do not have marked effects on growth (Mosher et al. 2009).

Pollen expresses both 21-nt and 24-nt siRNAs (Calarco et al. 2012); 21-nt siRNAs mediate posttranscriptional repression (Mosher 2010) and might target matrigenic mRNAs in early endosperm. And 24-nt siRNAs target the promoters of MEGs in pollen and possibly contribute to the silencing of paternal alleles in endosperm (Calarco et al. 2012). Unlike the maternally expressed siRNAs that are conjectured to target growth enhancers, paternally expressed siRNAs that target growth inhibitors are not expressed in postfertilization endosperm. This makes

sense. Postfertilization expression of paternal siRNAs would result in inactivation of maternal growth-inhibitor alleles in self-fertilized (or otherwise homozygous) endosperms, whereas the strategic patrigenic response to homozygosity is greater growth inhibition.

Evolution of Imprinted siRNAs

A *trans*-acting siRNA matches, but is unlinked to, its target, whereas a *cum*-acting siRNA is directly encoded by its target. Both kinds of maternally expressed siRNAs are predicted to target growth enhancers. The initial match of a *trans*-acting siRNA to a target is fortuitous, with natural selection sifting siRNAs with appropriate targets from siRNAs with inappropriate targets, whereas a *cum*-acting siRNA necessarily matches itself. siRNA-mediated inhibition may be evolutionarily stable if a *trans*-acting siRNA matches a functionally conserved region of its target gene, but *trans*-acting siRNAs that match genic regions that can evolve to evade the match (e.g., via synonymous base changes) will have evolutionarily transient effects.

Imprinted siRNAs that target their host gene have unusual evolutionary properties. This section can only sketch the complex interactions among alleles, and its tentative conclusions will need to be validated with more formal models. Consider two alleles at a growth-enhancing locus of an outcrossing plant: A' contains a siRNA-generating sequence (siDNA) and is sensitive to the encoded siRNA; A lacks siDNA and is insensitive to the siRNA of A' . The alleles are otherwise equivalent, with the expression of A being an evolutionary compromise between a lower level favored as a matrigene and a higher level favored as a patrigene. A' , when rare, will usually be inherited from one parent, not both, and will be expressed at higher levels as a patrigene than as a matrigene, because it is only in the latter role that its siRNA is expressed (**Figure 4**). Therefore, A' will increase in frequency because it imposes lesser demands on mothers as a matrigene.

Once A' increases in frequency, endosperms will often inherit A' from both parents. As a result, patrigenic A' encounters siRNA transcribed from matrigenic A' , and both alleles are repressed. Patrigenic A' learns that its seed contains an $A'A'$ embryo rather than an AA' embryo (expected relatedness to its own embryo is doubled) and that at least 50% of sibling embryos on the maternal sporophyte carry matrigenic A' (expected relatedness to other embryos more than doubled). Therefore, the optimal trade-off for patrigenic A' shifts toward smaller seeds and more efficient use of maternal resources. Strategic use of this information would favor less production of growth enhancer, precisely the effect of the siRNA.

A' performs a neat trick. It is associated with highest levels of growth enhancer when it is a patrigene in single dose (AAA'), intermediate levels when a matrigene in double dose ($A'A'A$), and lowest levels in triple dose ($A'A'A'$). Triploid endosperm may provide favorable stoichiometry for matrigenic silencing of patrigenes. In an $A'A'$ diploid endosperm, siRNA transcribed from a single allele would have two targets to silence, whereas, in an $A'A'A'$ triploid endosperm, siRNA transcribed from two alleles has three targets to silence.

Repression of patrigenic A' in $A'A'A'$ endosperms is a serendipitous effect that kicks in as A' increases in frequency in outcrossing populations. By contrast, in plants that can self-fertilize, selfed seeds will contain $A'A'A'$ endosperms, even when A' is rare. The siRNA causes patrigenic A' to be expressed at lower levels in self-fertilized than in outcrossed seeds, which increases the genic fitness of A' and individual fitness of mothers (De Jong et al. 2005).

The growth-inhibitory effects of A' are likely to be evolutionarily transient because selection favors replacement of A' by A'' , an allele that retains the siRNA but is relatively insensitive to its effects. A'' shifts expression toward the level before invasion of A' . In a sense, nothing has changed, except the genome has acquired an additional siRNA to which the host gene is relatively insensitive (**Figure 4**). The system is primed for the introduction of another siRNA. This iterative




Figure 4

The small-interfering RNA (siRNA) ratchet at a growth-enhancing locus. A population fixed for an unimprinted allele *A* (red, top left) is invaded by *A'*, a version of *A* that incorporates a maternally expressed siRNA (green, center). *A'* is in turn displaced by *A''*, a version of *A'* that is relatively insensitive to its own siRNA (purple, bottom right). The triploid endosperm contains two maternal alleles (upper and middle alleles) and one paternal allele (lower allele). The main diagonal represents homozygous endosperms, and the off-diagonal elements represent heterozygous endosperms.

process could explain the great diversity of maternally expressed siRNAs, their rapid evolutionary turnover, and the mildness of their effects (Ma et al. 2010, Mosher et al. 2009).

The imprinted siRNA acts as a green beard when *A* and *A'* are the only alleles, causing patrigenic *A'* to be less demanding in the presence of matrigenic *A'*. The siRNA ceases to be a reliable marker of self once *A''* also produces siRNA. Matrigenic *A''* induces patrigenic *A'* to reduce demand but does not reciprocate when matrigenic and patrigenic roles are reversed.

Previous models of the evolution of imprinted expression have considered alleles with expression x_m as a matrigene and x_p as a patrigene. These models found that the unbeatable strategy at a growth-enhancing locus is $x_m = 0$, $x_p = X_p$, where X_p is the level of expression optimal for a rare allele in its patrigenic role (Haig 1997, Wilkins & Haig 2001). The models assumed x_m and x_p were constant properties of an allele, determined in *cis*. Effects in *cum* violate this assumption and allow sophisticated strategies in which an allele changes its expression conditional on the identity of the other allele in its nucleus, and in which one allele can trick its partner to reduce expression. This richer set of strategies appears to facilitate lower levels of overt conflict, with production of growth enhancer maintained below X_p .

ORIGIN AND SIGNIFICANCE OF ENDOSPERM

Segregation of parental alleles at meiosis and sharing of offspring by mothers and fathers create genetic conflicts over the optimal size of seeds. Maternal genes are predicted to favor the smallest

seeds but the greatest seed number, with matrigenes then patrigenes favoring progressively larger but fewer seeds. The expected number of surviving offspring declines as seed size increases above the maternal optimum (Haig 1992).

Triploid endosperm may allow greater maternal control of the distribution of resources among seeds (Westoby & Rice 1982). Matrigenes of endosperm have a couple of intrinsic advantages over patrigenes. First, female gametophytes contribute more cytoplasm to early endosperm than do male gametophytes. Second, each matrigenes is present in two copies for each copy of a patrigene. Matrigenes are therefore expected to exert more phenotypic power than do patrigenes in endosperm. Maternally expressed siRNAs may allow seed size to be maintained closer to the maternal optimum than would be otherwise possible. Growth-enhancing genes that incorporate imprinted siRNAs are predicted to exhibit imprinted expression when heterozygous but unimprinted expression when homozygous and are predicted to be expressed at lower levels than conventionally imprinted genes would be.

The twin sperms of *Arabidopsis* appear to be functionally interchangeable (Ingouff et al. 2009). It seems reasonable to suppose that the sperm nucleus that first fertilized proto-endosperm of ancestral angiosperms contained the same epigenetic signature as the sperm nucleus that fertilized the egg. By contrast, egg and central-cell nuclei of modern angiosperms are epigenetically distinct (Moll et al. 2008). The much-debated question of the gametophytic versus embryonic origin of endosperm (Friedman 2001) may miss the mark if proto-endosperm was paternally embryonic but maternally gametophytic.

FUTURE ISSUES

1. Considerable progress has been made in understanding the genetic control of seed size once a commitment is made to provisioning a seed. However, many plants produce, and have fertilized, many more ovules than they can provision. Relatively little is known about the genetic control of adaptive seed abortion.
2. The evolution of siRNAs affecting seed size is expected to differ for siRNAs that target their own host sequence (*cum* effect) and those that target sequences at a different locus (*trans* effect). The relative importance of *cum* and *trans* effects must be addressed.
3. siRNAs with *cum* effects are subject to different selective forces in self-fertilizing and outcrossing plants. It will be important to determine whether observations in self-fertilizing *Arabidopsis thaliana* are also typical of outcrossing plants.
4. The identification of the targets of imprinted siRNAs will test their proposed role in kin conflict.

DISCLOSURE STATEMENT

The author is not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

The review has benefited from the comments of Jeffrey Chen, Kathleen Coleman, Rebecca Mosher, and Robert Trivers.

LITERATURE CITED

- Adamski NM, Anastasiou E, Eriksson S, O'Neill CM, Lenhard M. 2009. Local maternal control of seed size by *KLUH/CYP78A5*-dependent growth signaling. *Proc. Natl. Acad. Sci. USA* 106:20115–20
- Alonso-Blanco C, Blankestijn-de Vries H, Hanhart CJ, Kornneef M. 1999. Natural allelic variation at seed size loci in relation to other life history traits in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 96:4710–17
- Autran D, Baroux C, Raissig MT, Lenormand T, Wittig M, et al. 2011. Maternal epigenetic pathways control parental contributions to *Arabidopsis* early embryogenesis. *Cell* 145:707–19
- Baroux C, Gagliardini V, Page DR, Grossniklaus U. 2006. Dynamic regulatory interactions of *Polycomb* group genes: *MEDEA* autoregulation is required for imprinted gene expression in *Arabidopsis*. *Genes Dev.* 20:1081–86
- Baud S, Boutin JP, Miquel M, Lepiniec L, Rochat C. 2002. An integrated overview of seed development in *Arabidopsis thaliana* ecotype WS. *Plant Physiol. Biochem.* 40:151–60
- Bayer M, Nawy T, Giglione C, Galli M, Meinel T, Lukowitz W. 2009. Paternal control of embryonic patterning in *Arabidopsis thaliana*. *Science* 323:1485–88
- Berger F, Grini PE, Schnittger A. 2006. Endosperm: an integrator of seed growth and development. *Curr. Opin. Plant Biol.* 9:664–70
- Brandvain Y, Haig D. 2005. Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. *Am. Nat.* 166:330–38**
- Brown RC, Lemmon BE, Nguyen H, Olsen OA. 1999. Development of endosperm in *Arabidopsis thaliana*. *Sex. Plant Reprod.* 12:32–42
- Burt A, Trivers R. 1998. Genetic conflicts in genomic imprinting. *Proc. R. Soc. B* 265:2393–97
- Bushell C, Spielman M, Scott RJ. 2003. The basis of natural and artificial postzygotic hybridization barriers in *Arabidopsis* species. *Plant Cell* 15:1430–42
- Calarco JP, Borges F, Donoghue MTA, Van Ex F, Jullien PE, et al. 2012. Reprogramming of DNA methylation in pollen guides epigenetic inheritance via small RNA. *Cell* 151:194–204
- Charlesworth B, Langley CH. 1986. The evolution of self-regulated transposition of transposable elements. *Genetics* 112:359–83
- Charlton WL, Keen CL, Merriman C, Lynch P, Greenland AJ, Dickinson HG. 1995. Endosperm development in *Zea mays*; implications of gametic imprinting and paternal excess in regulation of transfer layer development. *Development* 121:3089–97
- Chaudhury AM, Ming L, Miller C, Craig S, Dennis ES, Peacock WJ. 1997. Fertilization-independent seed development in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 94:4223–28
- Chen LJ, Diao ZY, Specht C, Sung ZR. 2009. Molecular evolution of *VEF*-domain-containing *PcG* genes in plants. *Mol. Plant* 2:738–54
- Choi Y, Gehring M, Johnson L, Hannon M, Harada JJ, et al. 2002. DEMETER, a DNA glycosylase domain protein, is required for endosperm gene imprinting and seed viability in *Arabidopsis*. *Cell* 110:33–42
- Costa LM, Gutiérrez-Marcos JF, Brutnell TP, Greenland AJ, Dickinson HG. 2003. The *globby1-1* (*glo1-1*) mutation disrupts nuclear and cell division in the developing maize seed causing alterations in endosperm cell fate and tissue differentiation. *Development* 130:5009–17
- Costa LM, Yuan Y, Rouster J, Paul W, Dickinson H, Gutiérrez-Marcos JF. 2012. Maternal control of nutrient allocation in plant seeds by genomic imprinting. *Curr. Biol.* 22:160–65
- Curley JP, Barton S, Surani A, Keverne EB. 2004. Coadaptation in mother and infant regulated by a paternally expressed imprinted gene. *Proc. R. Soc. B* 271:1303–9
- Curtis MD, Grossniklaus U. 2008. Molecular control of autonomous embryo and endosperm development. *Sex. Plant Reprod.* 21:79–88
- Danilevskaya ON, Hermon P, Hantke S, Muszynski MG, Kollipara K, et al. 2003. Duplicated *fie* genes in maize: expression pattern and imprinting suggest distinct functions. *Plant Cell* 15:425–38
- De Jong TJ, Hermans CM, Van Der Veen-van Wijk KCAM. 2011. Paternal effects on seed mass in *Arabidopsis thaliana*. *Plant Biol.* 13(Suppl. 1):71–77
- De Jong TJ, Van Dijk H, Klinkhamer PGL. 2005. Hamilton's rule, imprinting and parent-offspring conflict over seed mass in partially selfing plants. *J. Evol. Biol.* 18:676–82

Describes how changes in mating system help to explain postzygotic incompatibilities between species.

- Dickinson H, Costa L, Gutierrez-Marcos J. 2012. Epigenetic neofunctionalisation and regulatory gene evolution in grasses. *Trends Plant Sci.* 17:389–94
- Dilkes BP, Comai L. 2004. A differential dosage hypothesis for parental effects in seed development. *Plant Cell* 16:3174–80
- Dilkes BP, Spielman M, Weizbauer R, Watson B, Burkart-Waco D, et al. 2008. The maternally expressed WRKY transcription factor *TTG2* controls lethality in interploidy crosses of *Arabidopsis*. *PLoS Biol.* 6:e308
- FitzGerald J, Luo M, Chaudhury A, Berger F. 2008. DNA methylation causes predominant maternal controls of plant embryo growth. *PLoS ONE* 3:e2298
- FitzGerald JN, Hui PS, Berger F. 2009. Polycomb group-dependent imprinting of the actin regulator *AtFH5* regulates morphogenesis in *Arabidopsis thaliana*. *Development* 136:3399–404
- Friedman WE. 2001. Developmental and evolutionary hypotheses for the origin of double fertilization and endosperm. *C. R. Acad. Sci. Paris* 324:559–67
- Garcia D, FitzGerald JN, Berger F. 2005. Maternal control of integument cell elongation and zygotic control of endosperm growth are coordinated to determine seed size in *Arabidopsis*. *Plant Cell* 17:52–60**
- Garcia D, Saingery V, Chambrier P, Mayer U, Jürgens G, Berger F. 2003. *Arabidopsis haiku* mutants reveal new controls of seed size by endosperm. *Plant Physiol.* 131:1661–70
- Gehring M, Bubb KL, Henikoff S. 2009. Extensive demethylation of repetitive elements during seed development underlies gene imprinting. *Science* 324:1447–51
- Gehring M, Huh JH, Hsieh T-F, Choi Y, Harada JJ, et al. 2006. DEMETER DNA glycosylase establishes *MEDEA* Polycomb gene self-imprinting by allele-specific demethylation. *Cell* 124:495–506
- Gehring M, Missirian V, Henikoff S. 2011. Genomic analysis of parent-of-origin allelic expression in *Arabidopsis thaliana* seeds. *PLoS ONE* 6:e23687
- Godfray HCJ. 1995. Evolutionary theory of parent-offspring conflict. *Nature* 376:133–38
- Grossniklaus U, Vielle-Calzada JP, Hoepfner MA, Gagliano WB. 1998. Maternal control of embryogenesis by *MEDEA*, a Polycomb group gene in *Arabidopsis*. *Science* 280:446–50
- Guitton AE, Berger F. 2011. Control of reproduction by Polycomb Group complexes in animals and plants. *Int. J. Dev. Biol.* 49:707–16
- Gutiérrez-Marcos JF, Costa LM, Biderre-Petit C, Khbaya B, O’Sullivan DM, et al. 2004. *maternally expressed gene1* is a novel maize endosperm transfer cell-specific gene with a maternal parent-of-origin pattern of expression. *Plant Cell* 16:1288–301
- Gutiérrez-Marcos JF, Pennington PD, Costa LM, Dickinson HG. 2003. Imprinting in the endosperm: a possible role in preventing wide hybridization. *Phil. Trans. R. Soc. B* 358:1105–11
- Haig D. 1992. Genomic imprinting and the theory of parent-offspring conflict. *Semin. Dev. Biol.* 3:153–60
- Haig D. 1996. Gestational drive and the green-bearded placenta. *Proc. Natl. Acad. Sci. USA* 93:6547–51
- Haig D. 1997. Parental antagonism, relatedness asymmetries, and genomic imprinting. *Proc. R. Soc. B* 264:1657–62
- Haig D. 2012. Retroviruses and the placenta. *Curr. Biol.* 22:R609–13
- Haig D. 2013. Parent-offspring coadaptation and conflict in the evolution of genomic imprinting. *Heredity* 275:1823–30
- Haig D, Trivers R. 1995. The evolution of parental imprinting: a review of hypotheses. In *Genomic Imprinting: Causes and Consequences*, ed. R Ohlsson, K Hall, M Ritzen, pp. 17–28. Cambridge: Cambridge Univ. Press
- Haig D, Westoby M. 1988. Inclusive fitness, seed resources and maternal care. In *Plant Reproductive Ecology*, ed. J Lovett-Doust, L Lovett-Doust, pp. 60–79. New York: Oxford Univ. Press
- Haig D, Westoby M. 1989. Parent-specific gene expression and the triploid endosperm. *Am. Nat.* 134:147–55**
- Haig D, Westoby M. 1991. Genomic imprinting in endosperm: its effects on seed development in crosses between species and between different ploidy levels of the same species, and its implications for the evolution of apomixis. *Phil. Trans. R. Soc. B* 333:1–13
- Hollick JB. 2012. Paramutation: a *trans*-homolog interaction affecting heritable gene regulation. *Curr. Opin. Plant Biol.* 15:536–43
- Hsieh TF, Ibarra CA, Silva P, Zemach A, Eshed-Williams L, et al. 2009. Genome-wide demethylation of *Arabidopsis* endosperm. *Science* 324:1451–54

Explains how seed size in *Arabidopsis* is determined by the interaction between maternal control of integument growth and filial control of endosperm growth.

First presented the parental conflict theory of the evolution of genomic imprinting.

- Hsieh TF, Shin J, Uzawa R, Silva P, Cohen S, et al. 2011. Regulation of imprinted gene expression in *Arabidopsis* endosperm. *Proc. Natl. Acad. Sci. USA* 108:1755–62
- Hughes R, Spielman M, Schruff MC, Larson TR, Graham IA, Scott RJ. 2008. Yield assessment of integument-led seed growth following targeted repair of *auxin response factor 2*. *Plant Biotechnol. J.* 6:758–69
- Ikeda Y. 2012. Plant imprinted genes identified by genome-wide approaches and their regulatory mechanisms. *Plant Cell Physiol.* 53:809–16
- Ingouff M, FitzGerald JN, Guérin C, Robert H, Sørensen MB, et al. 2005a. Plant formin AtFH5 is an evolutionarily conserved actin nucleator involved in cytokinesis. *Nat. Cell Biol.* 7:374–80
- Ingouff M, Haseloff J, Berger F. 2005b. Polycomb group genes control developmental timing of endosperm. *Plant J.* 42:663–74
- Ingouff M, Sakata T, Li J, Sprunck S, Dresselhaus T, Berger F. 2009. The two male gametes share equal ability to fertilize the egg cell in *Arabidopsis thaliana*. *Curr. Biol.* 19:R19–20
- Jahnke S, Scholten S. 2009. Epigenetic resetting of a gene imprinted in plant embryos. *Curr. Biol.* 19:1677–81
- Jiang H, Köhler C. 2012. Evolution, function, and regulation of genomic imprinting in plant seed development. *J. Exp. Bot.* 63:4713–22
- Johnston SA, den Nijs TPM, Peloquin SJ, Hanneman RE. 1980. The significance of genic balance to endosperm development in interspecific crosses. *Theor. Appl. Genet.* 57:5–9
- Johnston SA, Hanneman RE. 1982. Manipulations of endosperm balance number overcome crossing barriers between diploid *Solanum* species. *Science* 217:446–48
- Josefsson C, Dilkes B, Comai L. 2006. Parent-dependent loss of gene silencing during interspecies hybridization. *Curr. Biol.* 16:1322–28
- Jullien PE, Berger F. 2010. Parental genome dosage imbalance deregulates imprinting in *Arabidopsis*. *PLoS Genet.* 6:e1000886
- Jullien PE, Katz A, Oliva M, Ohad N, Berger F. 2006. Polycomb group complexes self-regulate imprinting of the Polycomb group gene *MEDEA* in *Arabidopsis*. *Curr. Biol.* 16:486–92
- Jullien PE, Susaki D, Yelagandula R, Higshiyama T, Berger F. 2012. DNA dynamics during sexual reproduction in *Arabidopsis thaliana*. *Curr. Biol.* 22:1825–30
- Kang B-H, Xiong Y, Williams DS, Pozueta-Romero D, Chourey PS. 2010. *Miniature1*-encoded cell wall invertase is essential for assembly and function of wall-in-growth in the maize endosperm transfer cell. *Plant Physiol.* 151:1366–76
- Kang I-H, Steffen JG, Portereiko MF, Lloyd A, Drews GN. 2008. The AGL62 MADS domain protein regulates cellularization during endosperm development in *Arabidopsis*. *Plant Cell* 20:635–47
- Kawabe A, Fujimoto R, Charlesworth D. 2007. High diversity due to balancing selection in the promoter region of the *Medea* gene in *Arabidopsis lyrata*. *Curr. Biol.* 17:1885–89
- Keverne EB, Curley JP. 2008. Epigenetics, brain evolution and behaviour. *Front. Neuroendocrinol.* 29:398–412
- Kinoshita T, Miura A, Choi Y, Kinoshita Y, Cao X, et al. 2004. One-way control of *FWA* imprinting in *Arabidopsis* endosperm by DNA methylation. *Science* 303:521–23
- Kiyosue T, Ohad N, Yadegari R, Hannon M, Dinneny J, et al. 1999. Control of fertilization-independent seed development by the *MEDEA* polycomb gene in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 96:4186–91
- Köhler C, Kradolfer D. 2011. Epigenetic mechanisms in the endosperm and their consequences for the evolution of flowering plants. *Biochim. Biophys. Acta* 1809:438–43
- Köhler C, Makarevich G. 2006. Epigenetic mechanisms governing seed development in plants. *EMBO Rep.* 7:1223–27
- Köhler C, Weinhofer-Molisch I. 2010. Mechanisms and evolution of genomic imprinting in plants. *Heredity* 105:57–63
- Köhler C, Hennig L, Spillane C, Pien S, Grussem W, Grossniklaus U. 2003. The *Polycomb*-group protein *MEDEA* regulates seed development by controlling expression of the MADS-box gene *PHERES1*. *Genes Dev.* 17:1540–53
- Köhler C, Page DR, Gagliardini V, Grossniklaus U. 2005. The *Arabidopsis thaliana* *MEDEA* Polycomb group protein controls expression of *PHERES1* by parental imprinting. *Nat. Genet.* 37:28–30
- Köhler C, Wolff P, Spillane C. 2012. Epigenetic mechanisms underlying genomic imprinting in plants. *Annu. Rev. Plant Biol.* 63:331–52

- Kondoh M, Higashi M. 2000. Reproductive isolation mechanisms resulting from resolution of intragenomic conflict. *Am. Nat.* 156:511–18
- Lauria M, Rupe M, Guo M, Kranz E, Pirona R, et al. 2004. Extensive maternal hypomethylation in the endosperm of *Zea mays*. *Plant Cell* 16:510–22
- Leblanc O, Pointe C, Hernandez M. 2002. Cell cycle progression during endosperm development in *Zea mays* depends on parental dosage effects. *Plant J.* 32:1057–66
- Li J, Berger F. 2012. Endosperm: food for mankind and fodder for scientific discoveries. *New Phytol.* 195:290–305
- Li N, Dickinson HG. 2010. Balance between maternal and paternal alleles sets the timing of resource accumulation in the maize endosperm. *Proc. R. Soc. B* 277:3–10
- Li Y, Zheng L, Corke F, Smith C, Bevan MW. 2008. Control of final seed and organ size by the *DA1* gene family in *Arabidopsis thaliana*. *Genes Dev.* 22:1331–36
- Lin BY. 1984. Ploidy barrier to endosperm development in maize. *Genetics* 107:103–15**
- Lippman Z, Gendrek AV, Black M, Vaughn MW, Dedhia N, et al. 2004. Role of transposable elements in heterochromatin and epigenetic control. *Nature* 430:471–76
- Lisch D, Slotkin RK. 2011. Strategies for silencing and escape: the ancient struggle between transposable elements and their hosts. *Int. Rev. Cell Mol. Biol.* 292:119–52
- Liu S-L, Adams KL. 2010. Dramatic change in function and expression pattern of a gene duplicated by polyploidy created a paternal effect gene in the Brassicaceae. *Mol. Biol. Evol.* 27:2817–28
- Lu J, Zhang C, Baulcombe DC, Chen ZJ. 2012. Maternal siRNAs as regulators of parental genome imbalance and gene expression in endosperm of *Arabidopsis* seeds. *Proc. Natl. Acad. Sci. USA* 109:5529–34**
- Luo M, Bilodeau P, Dennis ES, Peacock WJ, Chaudhury A. 2000. Expression and parent-of-origin effects for *FIS2*, *MEA*, and *FIE* in the endosperm and embryo of developing *Arabidopsis* seeds. *Proc. Natl. Acad. Sci. USA* 97:10637–42
- Luo M, Bilodeau P, Koltunow A, Dennis ES, Peacock WJ, Chaudhury AM. 1999. Genes controlling fertilization-independent seed development in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 96:296–301
- Luo M, Dennis ES, Berger F, Peacock WJ, Chaudhury A. 2005. *MINISEED3 (MINI3)*, a WRKY family gene, and *HAIKU2 (IKU2)*, a leucine-rich repeat (*LRR*) *KINASE* gene are regulators of seed size in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 102:17531–36
- Luo M, Platten D, Chaudhury A, Peacock WJ, Dennis ES. 2009. Expression, imprinting, and evolution of rice homologs of the polycomb group genes. *Mol. Plant* 2:711–23
- Luo M, Taylor JM, Spriggs A, Zhang H, Wu X, et al. 2011. A genome-wide survey of imprinted genes in rice seeds reveals imprinting primarily occurs in the endosperm. *PLoS Genet.* 6:e1002125
- Ma Z, Coruh C, Axtell MJ. 2010. *Arabidopsis lyrata* small RNAs: transient *MIRNA* and small interfering RNA loci within the *Arabidopsis* genus. *Plant Cell* 22:1090–103
- Makarevich G, Leroy O, Akinci U, Schubert D, Clarenz O, et al. 2006. Different *Polycomb* group complexes regulate common target genes in *Arabidopsis*. *EMBO Rep.* 7:947–52
- Makarevich G, Villar CBR, Erilova A, Köhler C. 2008. Mechanism of *PHERES1* imprinting in *Arabidopsis*. *J. Cell Sci.* 121:906–12
- Mansfield SG, Briarty LG. 1991. Early embryogenesis in *Arabidopsis thaliana*. II. The developing embryo. *Can. J. Bot.* 69:461–76
- Mansfield SG, Briarty LG. 1992. Cotyledon cell development in *Arabidopsis thaliana* during reserve deposition. *Can. J. Bot.* 70:151–64
- Martienssen RA. 2010. Heterochromatin, small RNA and post-fertilization dysgenesis in allopolyploid and interloid hybrids of *Arabidopsis*. *New Phytol.* 186:46–53
- Matzke MA, Matzke AJM, Eggleston WB. 1996. Paramutation and transgene silencing: A common response to invasive DNA? *Trends Plant Sci.* 1:382–88
- McVean GT, Hurst LD. 1997. Molecular evolution of imprinted genes: no evidence for antagonistic coevolution. *Proc. R. Soc. B* 264:739–46
- Miyake T, Takebayashi N, Wolf DE. 2009. Possible diversifying selection in the imprinted gene *MEDEA* in *Arabidopsis*. *Mol. Biol. Evol.* 26:843–57

Definitively demonstrated that normal endosperm development depended on a 2m:1p ratio of parental genomes.

Describes how maternally expressed siRNAs inhibit the proliferation of endosperm.

First demonstrated
imprinted siRNAs.

- Moll C, Nielsen N, Groß-Hardt R. 2008. Mutants with aberrant numbers of gametic cells shed new light on old questions. *Plant Biol.* 10:529–33
- Molnar A, Melnyk CW, Bassett A, Hardcastle TJ, Dunn R, Baulcombe DC. 2010. Small silencing RNAs in plants are mobile and direct epigenetic modification in recipient cells. *Science* 328:872–75
- Morley-Smith ER, Pike MJ, Findlay K, Köckenberger W, Hill LM, et al. 2008. The transport of sugars to developing embryos is not via the bulk endosperm in oilseed rape seeds. *Plant Physiol.* 147:2121–30
- Mosher RA. 2010. Maternal control of Pol IV-dependent siRNAs in *Arabidopsis* endosperm. *New Phytol.* 186:358–64
- Mosher RA, Melnyk CW. 2010. siRNAs and DNA methylation: seedy epigenetics. *Trends Plant Sci.* 15:204–10
- Mosher RA, Melnyk CW, Kelly KA, Dunn RM, Studholme DJ, Baulcombe DC. 2009. Uniparental expression of PolIV-dependent siRNAs in developing endosperm of *Arabidopsis*. *Nature* 460:283–86**
- Mosher RA, Tan EH, Shin J, Fischer RL, Pikaard CS, Baulcombe DC. 2011. An atypical epigenetic mechanism affects uniparental expression of Pol IV-dependent siRNAs. *PLoS ONE* 6:e25756
- Nishiyama I, Yabuno T. 1978. Causal relationships between the polar nuclei in double fertilization and interspecific cross-incompatibility in *Avena*. *Cytologia* 43:453–66
- Nishiyama I, Yabuno T. 1979. Triple fusion of the primary endosperm nucleus as a cause of interspecific cross-incompatibility in *Avena*. *Euphytica* 28:57–65
- Nodine MD, Bartel DP. 2012. Maternal and paternal genomes contribute equally to the transcriptome of early plant embryos. *Nature* 482:94–97
- Ohto M, Floyd SK, Fischer RL, Goldberg RB, Harada JJ. 2009. Effects of APETALA2 on embryo, endosperm, and seed coat development determine seed size in *Arabidopsis*. *Sex. Plant Reprod.* 22:277–89
- Olsen OA. 2004. Nuclear endosperm development in cereals and *Arabidopsis thaliana*. *Plant Cell* 16(Suppl.):S214–27
- Pennington PD, Costa LM, Gutiérrez-Marcos JF, Greenland AJ, Dickinson HG. 2008. When genomes collide: aberrant seed development following maize interploidy crosses. *Ann. Bot.* 101:833–43
- Pignatta D, Gehring M. 2012. Imprinting meets genomics: new insights and new challenges. *Curr. Opin. Plant Biol.* 15:530–35
- Queller DC. 1983. Kin selection and conflict in seed maturation. *J. Theor. Biol.* 100:153–72**
- Queller DC. 1989. Inclusive fitness in a nutshell. *Oxf. Surv. Evol. Biol.* 6:73–109
- Queller DC. 2003. Theory of genomic imprinting conflict in social insects. *BMC Evol. Biol.* 3:15
- Rodrigues JCM, Luo M, Berger F, Koltunow AMG. 2010. Polycomb group gene function in sexual and asexual seed development in angiosperms. *Sex. Plant Reprod.* 22:123–33
- Schoft VK, Chumak N, Choi Y, Hannon M, Garcia-Aguilar M, et al. 2011. Function of the DEMETER DNA glycosylase in the *Arabidopsis thaliana* male gametophyte. *Proc. Natl. Acad. Sci. USA* 108:8042–47
- Schruff MC, Spielman M, Tiwari S, Adams S, Fenby N, Scott RJ. 2006. The *AUXIN RESPONSE FACTOR 2* gene of *Arabidopsis* links auxin signalling, cell division, and the size of seeds and other organs. *Development* 133:251–61
- Schubert D, Goodrich J. 2003. Plant epigenetics: MEDEA's children take centre stage. *Curr. Biol.* 13:R638–40
- Scott RJ, Spielman M, Bailey J, Dickinson HG. 1998. Parent-of-origin effects on seed development in *Arabidopsis thaliana*. *Development* 125:3329–41
- Shirzadi R, Andersen ED, Bjerkan KN, Gloekle BM, Heese M, et al. 2011. Genome-wide transcript profiling of endosperm without paternal contribution identifies parent-of-origin-dependent regulation of *AGAMOUS-LIKE36*. *PLoS Genet.* 7:e1001303
- Slotkin RK, Martienssen R. 2007. Transposable elements and the epigenetic regulation of the genome. *Nat. Rev. Genet.* 8:272–85
- Slotkin RK, Vaughn M, Borges F, Tanurdzi M, Becker JD, et al. 2009. Epigenetic reprogramming and small RNA silencing of transposable elements in pollen. *Cell* 136:461–72
- Smith CC, Fretwell SD. 1974. The optimal balance between size and number of offspring. *Am. Nat.* 108:499–506
- Spillane C, Schmid KJ, Laouecillé-Duprat S, Pien S, Escobar-Restrepo JM, et al. 2007. Positive darwinian selection at the imprinted *MEDEA* locus in plants. *Nature* 448:349–52

Represents an
important early
discussion of kin
conflicts during seed
development.

- Springer NM, Danilevskaia ON, Hermon P, Helentjaris TG, Phillips RL, et al. 2002. Sequence relationships, conserved domains, and expression patterns for maize homologs of the Polycomb group genes *E(z)*, *esc*, and *E(pc)*. *Plant Physiol.* 128:1332–45
- Stoute AI, Varenko V, King GJ, Scott RJ, Kurup S. 2012. Parental genome imbalance in *Brassica oleracea* causes asymmetric triploid block. *Plant J.* 71:503–16
- Swaney WT, Curley JP, Champagne FA, Keverne EB. 2007. Genomic imprinting mediates sexual experience-dependent olfactory learning in male mice. *Proc. Natl. Acad. Sci. USA* 104:6084–89
- Trivers RL. 1974. Parent-offspring conflict. *Am. Zool.* 14:249–64
- Verona RI, Mann MRW, Bartolomei MS. 2003. Genomic imprinting: intricacies of epigenetic regulation in clusters. *Annu. Rev. Cell Dev. Biol.* 19:237–59
- Vielle-Calzada J-P, Thomas J, Spillane C, Coluccio A, Hoepfner MA, Grossniklaus U. 1999. Maintenance of genomic imprinting at the *Arabidopsis medea* locus requires zygotic *DDMI* activity. *Genes Dev.* 13:2971–82
- Vilhar B, Kladnik A, Blejec A, Chourey PS, Dermastia A. 2002. Cytometrical evidence that the loss of seed weight in the *miniature1* seed mutant of maize is associated with reduced mitotic activity in the developing endosperm. *Plant Physiol.* 129:23–30
- Villar CBR, Erilova A, Makarevich G, Trösch R, Köhler C. 2009. Control of *PHERES1* imprinting by direct tandem repeats. *Mol. Plant* 2:654–60
- Vinkenoog R, Spielman M, Adams S, Fischer RL, Dickinson HG, Scott RJ. 2000. Hypomethylation promotes autonomous endosperm development and rescues postfertilization lethality in *fie* mutants. *Plant Cell* 12:2271–82
- Walia H, Josefsson C, Dilkes B, Kirkbride R, Harada J, Comai L. 2009. Dosage-dependent deregulation of an AGAMOUS-like gene cluster contributes to interspecific incompatibility. *Curr. Biol.* 19:1128–32
- Wang A, Garcia D, Zhang H, Feng K, Chaudhury A, et al. 2010. The VQ motif protein IKU1 regulates endosperm growth and seed size in Arabidopsis. *Plant J.* 63:670–79
- Waters AJ, Makarevitch I, Eichten SR, Swanson-Wagner RA, Yeh CT, et al. 2011. Parent-of-origin effects on gene expression and DNA methylation in the maize endosperm. *Plant Cell* 23:4221–33
- West SA, Gardner A. 2010. Altruism, spite, and greenbeards. *Science* 327:1341–44
- Westoby M, Rice B. 1982. Evolution of the seed plants and inclusive fitness of plant tissues. *Evolution* 36:713–24**
- Wilkins JF, Haig D. 2001. Genomic imprinting of two antagonistic loci. *Proc. R. Soc. B* 268:1861–67
- Wöhrmann HJP, Gagliardini V, Raissig MT, Wehrle W, Arand J, et al. 2012. Identification of a DNA methylation-independent imprinting control region at the *Arabidopsis MEDEA* locus. *Genes Dev.* 26:1837–50
- Wolf JB, Hager R. 2006. A maternal-offspring coadaptation theory for the evolution of genomic imprinting. *PLoS Biol.* 4:e380
- Wolff P, Weinhofer I, Seguin J, Poszak P, Beisel C, et al. 2011. High-resolution analysis of parent-of-origin allelic expression in the *Arabidopsis* endosperm. *PLoS Genet.* 6:e1002126
- Wu SC, Zhang Y. 2012. Active DNA demethylation: Many roads lead to Rome. *Nat. Rev. Mol. Cell Biol.* 11:607–20
- Xiao W, Brown RC, Lemmon BE, Harada JJ, Goldberg RB, Fischer RL. 2006. Regulation of seed size by hypomethylation of maternal and paternal genomes. *Plant Physiol.* 142:1160–68
- Yadegari R, Kinoshita T, Lotan O, Cohen G, Katz A, et al. 2000. Mutations in the *FIE* and *MEA* genes that encode interacting polycomb proteins cause parent-of-origin effects on seed development by distinct mechanisms. *Plant Cell* 12:2367–81
- Zemach A, Kim MY, Silva P, Rodrigues JA, Dotson B, et al. 2010. Local DNA hypomethylation in rice endosperm. *Proc. Natl. Acad. Sci. USA* 107:18729–34
- Zhang M, Zhao H, Xie S, Chen J, Xu Y, et al. 2011. Extensive, clustered parental imprinting of protein-coding and noncoding RNAs in developing maize endosperm. *Proc. Natl. Acad. Sci. USA* 108:20042–47
- Zhou Y, Zhang X, Kang X, Zhao X, Zhang X, Ni M. 2009. SHORT HYPOCOTYL UNDER BLUE1 associates with *MINISEED3* and *HAIKU2* promoters in vivo to regulate *Arabidopsis* seed development. *Plant Cell* 21:106–17

Provides an important early discussion of kin conflicts during seed development.



Contents

Formation and Segmentation of the Vertebrate Body Axis <i>Bertrand Bénazéraf and Olivier Pourquié</i>	1
Mechanobiology and Developmental Control <i>Tadanori Mammoto, Akiko Mammoto, and Donald E. Ingber</i>	27
Mitogen-Activated Protein Kinase Pathways in Osteoblasts <i>Matthew B. Greenblatt, Jae-Hyuck Shim, and Laurie H. Glimcher</i>	63
Pancreas Organogenesis: From Lineage Determination to Morphogenesis <i>Hung Ping Shih, Allen Wang, and Maiké Sander</i>	81
Beyond the Niche: Tissue-Level Coordination of Stem Cell Dynamics <i>Lucy Erin O'Brien and David Bilder</i>	107
Hormonal Control of Stem Cell Systems <i>Dana Gancz and Lilach Gilboa</i>	137
Spermatogonial Stem Cell Self-Renewal and Development <i>Mito Kanatsu-Shinohara and Takashi Shinohara</i>	163
Kin Conflict in Seed Development: An Interdependent but Fractious Collective <i>David Haig</i>	189
microRNA Control of Mouse and Human Pluripotent Stem Cell Behavior <i>Tobias S. Greve, Robert L. Judson, and Robert Blelloch</i>	213
Something Silent This Way Forms: The Functional Organization of the Repressive Nuclear Compartment <i>Joan C. Ritland Politz, David Scalzo, and Mark Groudine</i>	241
Cytoskeletal Dynamics in <i>Caenorhabditis elegans</i> Axon Regeneration <i>Andrew D. Chisholm</i>	271

Integrative Mechanisms of Oriented Neuronal Migration in the Developing Brain <i>Irina Evsyukova, Charlotte Plestant, and E.S. Anton</i>	299
TRP Channels and Pain <i>David Julius</i>	355
Synaptic Laminae in the Visual System: Molecular Mechanisms Forming Layers of Perception <i>Herwig Baier</i>	385
Microtubule-Depolymerizing Kinesins <i>Claire E. Walczak, Sophia Gayek, and Ryoma Obi</i>	417
Kinesin-2: A Family of Heterotrimeric and Homodimeric Motors with Diverse Intracellular Transport Functions <i>Jonathan M. Scholey</i>	443
Microtubules in Cell Migration <i>Sandrine Etienne-Manneville</i>	471
Mathematical Modeling of Eukaryotic Cell Migration: Insights Beyond Experiments <i>Gaudenz Danuser, Jun Allard, and Alex Mogilner</i>	501
GTP-Dependent Membrane Fusion <i>James A. McNew, Holger Sondermann, Tina Lee, Mike Stern, and Federica Brandizzi</i>	529
Viral Membrane Scission <i>Jeremy S. Rossman and Robert A. Lamb</i>	551
How Microbiomes Influence Metazoan Development: Insights from History and <i>Drosophila</i> Modeling of Gut-Microbe Interactions <i>Won-Jae Lee and Paul T. Brey</i>	571
Cell and Developmental Biology of Arbuscular Mycorrhiza Symbiosis <i>Caroline Gutjahr and Martin Parniske</i>	593

Indexes

Cumulative Index of Contributing Authors, Volumes 25–29	619
Cumulative Index of Article Titles, Volumes 25–29	622

Errata

An online log of corrections to *Annual Review of Cell and Developmental Biology* articles may be found at <http://cellbio.annualreviews.org/errata.shtml>